

On an alternate proof of Hamilton's matrix Harnack inequality for the Ricci flow

Bennett Chow*

Department of Mathematics
University of California at San Diego

August 15, 2001

In [LY] a differential Harnack inequality was proved for solutions to the heat equation on a Riemannian manifold. Inspired by this result, Hamilton first proved trace and matrix Harnack inequalities for the Ricci flow on compact surfaces [H0] and then vastly generalized his own result to all higher dimensions for complete solutions of the Ricci flow with nonnegative curvature operator [H2]. Soon afterwards, a matrix Harnack inequality for the Kähler-Ricci flow under the assumption of nonnegative bisectional curvature was proved by Huai-Dong Cao [C]. In this paper, following a suggestion of Richard Hamilton, we give an alternate proof of the matrix Harnack inequality for the Ricci flow originally proved by him.¹ In [H2], Hamilton proved that if $(M^n, g(t))$ is a complete solution to the Ricci flow with nonnegative curvature operator, then

$$\frac{\partial R}{\partial t} + \frac{R}{t} + 2\nabla_i R V^i + 2R_{ij} V^i V^j \geq 0 \quad (1)$$

for any vector field V . In particular, if in addition the Ricci curvature is positive, then

$$\frac{\partial R}{\partial t} + \frac{R}{t} - \frac{1}{2} (Rc^{-1})^{ij} \nabla_i R \nabla_j R \geq 0.$$

This trace Harnack inequality is a corollary of Hamilton's matrix Harnack inequality, which we now recall (we follow Hamilton's sign convention for the curvature tensor so that $R_{ijkl} = g_{kh} R_{ijl}^h$). Define tensors P and M by

$$P_{abc} \doteq D_a R_{bc} - D_b R_{ac}$$

$$M_{ab} \doteq \Delta R_{ab} - \frac{1}{2} D_a D_b R + 2R_{acbd} R_{cd} - R_{ac} R_{bc} + \frac{1}{2t} R_{ab}$$

*Research partially supported by NSF grant DMS-9971891.

¹Yet another proof is given in [CC]. Hamilton's motivation for considering the alternate proof given in the present paper is that it may possibly facilitate extending his Harnack inequality to the variable signed curvature case, especially in dimension three. See the end of [H3] for some reasons for studying the problem of extending the Harnack inequality.

and consider the Harnack quadratic

$$Z(U, W) \doteq M_{ab}W_aW_b + 2P_{abc}U_{ab}W_c + R_{abcd}U_{ab}U_{cd},$$

where U is any 2-form and W is any 1-form. Hamilton's matrix Harnack estimate says that if $(M^n, g(t))$ is a complete solution to the Ricci flow with nonnegative curvature operator, then

$$Z(U, W) \geq 0$$

for all U and W . Taking $U = V \wedge W$ and summing over W in an orthonormal frame, one obtains (1).

The idea of giving an alternate proof of it is as follows. Given any vector W , consider the 2-form U minimizing the quadratic form. This leads to the consideration of a symmetric 2-tensor Z_{ab} which has a nice evolution equation. In particular, Z_{ab} is essentially a supersolution to the heat equation so that the maximum principle implies that it is positive definite, which is *equivalent* to the matrix Harnack inequality. Tracing this 2-tensor yields a trace Harnack inequality: $Z \doteq Z_{aa} > 0$.

Now we proceed with the details of the proof. Assume that the curvature operator is positive and let S_{abcd} denote the inverse of R_{abcd} so that

$$\sum_{e,f} S_{abef} R_{efcd} = I_{abcd},$$

where $I : \wedge^2 M \rightarrow \wedge^2 M$ is the identity. First we rewrite the matrix Harnack quadratic by completing the square:

$$\begin{aligned} Z(U, W) &= M_{ab}W_aW_b - S_{ijkl}P_{ijp}W_pP_{klq}W_q \\ &\quad + R_{abcd}(U_{ab} + S_{abij}P_{ijp}W_p)(U_{cd} + S_{cdkl}P_{klq}W_q). \end{aligned}$$

Since $Rm > 0$, the above expression is minimized when

$$U_{ab} + S_{abij}P_{ijp}W_p = 0,$$

in which case

$$Z(U, W) = (M_{ab} - S_{ijkl}P_{ija}P_{klb})W_aW_b.$$

Let

$$Z_{ab} \doteq M_{ab} - S_{ijkl}P_{ija}P_{klb},$$

which is a symmetric 2-tensor.

The main computation is that in an orthonormal moving frame, the evolution of Z_{ab} is given as follows. Following the idea in [H2], as long as it is nonnegative-

definite, we may write the Harnack quadratic as a sum of squares:

$$\begin{aligned} R_{abcd} &= \sum_N Y_{ab}^N Y_{cd}^N \\ P_{abc} &= \sum_N Y_{ab}^N X_c^N \\ M_{ab} &= \sum_N X_a^N X_b^N. \end{aligned}$$

Here, given a point $x \in M$, $\{Y^N \oplus X^N\}_{N=1}^m$ is some orthogonal set of vectors in $\wedge^2 M_x \oplus \wedge^1 M_x$ and $m = \dim \wedge^2 M_x + \dim \wedge^1 M_x = \frac{n(n+1)}{2}$.

Main Theorem. *If $Rm > 0$, then the Harnack quantity Z_{ab} satisfies the evolution equation:*

$$\left(\frac{\partial}{\partial t} - \Delta \right) Z_{ab} = 2S_{mntu} K_{avtu} K_{bvmn} + L_a^{NM} L_b^{NM} - \frac{2}{t} Z_{ab}, \quad (2)$$

where

$$\begin{aligned} K_{avtu} &\doteq S_{ijrs} P_{ija} \nabla_v R_{rstu} - \nabla_v P_{tua} + R_{tuaw} R_{vw} + \frac{1}{2t} R_{tuav} \\ L_a^{NM} &\doteq 2Y_{id}^N Y_{jd}^M S_{ijkl} P_{kla} + Y_{ah}^N X_h^M - Y_{ah}^M X_h^N. \end{aligned}$$

Note that applying the maximum principle for symmetric 2-tensors in section 7 of [H1] yields $Z_{ab} > 0$, which is equivalent to Hamilton's matrix Harnack inequality. Here we used the fact that if $Rm > 0$, then as long as Z_{ab} is nonnegative definite, L_a^{NM} is well-defined.

The bulk of the rest of this paper is devoted to proving the theorem. First recall Hamilton's computations for the evolution equations of the Harnack quadratic [H2].

Lemma 1 *In an evolving local orthonormal frame field, we have*

1.

$$(D_t - \Delta) R_{abcd} = 2(B_{abcd} - B_{abdc} + B_{acbd} - B_{adb c}), \quad (3)$$

where $B_{abcd} = R_{aebf} R_{cedf}$.

2.

$$(D_t - \Delta) P_{abc} = -2R_{de} D_d R_{abce} + 2R_{adbe} P_{dec} + 2R_{adce} P_{dbe} + 2R_{bdce} P_{ade}. \quad (4)$$

3.

$$\begin{aligned} (D_t - \Delta) M_{ab} &= 2R_{cd} [D_c P_{dab} + D_c P_{dba}] + 2R_{acbd} M_{cd} \\ &\quad + 2P_{acd} P_{bcd} - 4P_{acd} P_{bdc} + 2R_{cd} R_{ce} R_{adbe} - \frac{1}{2t^2} R_{ab}. \end{aligned} \quad (5)$$

We use this lemma in the computations below. We start by computing:

$$\begin{aligned} \left(\frac{\partial}{\partial t} - \Delta\right) Z_{ab} &= \left(\frac{\partial}{\partial t} - \Delta\right) M_{ab} - \left(\frac{\partial}{\partial t} - \Delta\right) S_{ijkl} P_{ija} P_{klb} \\ &\quad - S_{ijkl} \left(\frac{\partial}{\partial t} - \Delta\right) P_{ija} P_{klb} - S_{ijkl} P_{ija} \left(\frac{\partial}{\partial t} - \Delta\right) P_{klb} \\ &\quad + 2S_{ijkl} \nabla_p P_{ija} \nabla_p P_{klb} + 2\nabla_p S_{ijkl} \nabla_p P_{ija} P_{klb} + 2\nabla_p S_{ijkl} P_{ija} \nabla_p P_{klb}. \end{aligned}$$

Thus, besides the evolution equations for M_{ab} and P_{abc} , we need the evolution equation for S_{ijkl} , which is given by:

$$\begin{aligned} \left(\frac{\partial}{\partial t} - \Delta\right) S_{ijkl} &= -S_{ijmn} S_{klpq} \left(\frac{\partial}{\partial t} - \Delta\right) R_{mnpq} \\ &\quad - (S_{ijrs} S_{mntu} S_{klpq} + S_{ijmn} S_{klrs} S_{pqtu}) \nabla_v R_{rstu} \nabla_v R_{mnpq} \\ &= -2S_{ijmn} S_{klpq} (B_{mnpq} - B_{mnqp} + B_{mpnq} - B_{mqnp}) \\ &\quad - (S_{ijrs} S_{mntu} S_{klpq} + S_{ijmn} S_{klrs} S_{pqtu}) \nabla_v R_{rstu} \nabla_v R_{mnpq}. \end{aligned}$$

Substituting in those evolution equations, we obtain:

$$\begin{aligned} \left(\frac{\partial}{\partial t} - \Delta\right) Z_{ab} &= 2R_{cd} [D_c P_{dab} + D_c P_{dba}] + 2R_{acbd} M_{cd} \\ &\quad + 2P_{acd} P_{bcd} - 4P_{acd} P_{bdc} + 2R_{cd} R_{ce} R_{adbe} - \frac{1}{2t^2} R_{ab} \\ &\quad + \left((S_{ijrs} S_{mntu} S_{klpq} + S_{ijmn} S_{klrs} S_{pqtu}) \nabla_v R_{rstu} \nabla_v R_{mnpq} \right. \\ &\quad \left. + 2S_{ijmn} S_{klpq} (B_{mnpq} - B_{mnqp} + B_{mpnq} - B_{mqnp}) \right) P_{ija} P_{klb} \\ &\quad - 2S_{ijkl} P_{klb} (-R_{de} D_d R_{ijae} + R_{idje} P_{dea} + R_{idae} P_{dje} + R_{jdae} P_{ide}) \\ &\quad - 2S_{ijkl} P_{ija} (-R_{de} D_d R_{klbe} + R_{kdle} P_{deb} + R_{kdbe} P_{dle} + R_{ldbe} P_{kde}) \\ &\quad + 2S_{ijkl} \nabla_p P_{ija} \nabla_p P_{klb} - 2S_{ijmn} S_{klpq} \nabla_r R_{mnpq} (\nabla_r P_{ija} P_{klb} + P_{ija} \nabla_r P_{klb}) \end{aligned}$$

where $B_{abcd} = R_{aebf} R_{cedf}$. Taking into account symmetries and combining some like terms, we have

$$\begin{aligned} \left(\frac{\partial}{\partial t} - \Delta\right) Z_{ab} &= 2R_{acbd} Z_{cd} \\ &\quad + 2S_{ijkl} \nabla_p P_{ija} \nabla_p P_{klb} \\ &\quad - 2S_{ijmn} S_{klpq} \nabla_r R_{mnpq} (\nabla_r P_{ija} P_{klb} + P_{ija} \nabla_r P_{klb}) \\ &\quad + (S_{ijrs} S_{mntu} S_{klpq} + S_{ijmn} S_{klrs} S_{pqtu}) P_{ija} P_{klb} \nabla_v R_{rstu} \nabla_v R_{mnpq} \\ &\quad + 2R_{acbd} S_{ijkl} P_{ijc} P_{kld} + 2P_{acd} P_{bcd} - 4P_{acd} P_{bdc} \\ &\quad + 2S_{ijmn} S_{klpq} (B_{mnpq} - B_{mnqp} + B_{mpnq} - B_{mqnp}) P_{ija} P_{klb} \\ &\quad - 2(S_{dekl} R_{diej} + S_{ijde} R_{dkel}) P_{ija} P_{klb} \\ &\quad - 4S_{ijkl} R_{idae} P_{klb} P_{dje} - 4S_{ijkl} R_{kdbe} P_{ija} P_{dle} \\ &\quad + 2R_{cd} [D_c P_{dab} + D_c P_{dba}] + 2R_{cd} R_{ce} R_{adbe} - \frac{1}{2t^2} R_{ab} \\ &\quad + 2S_{ijkl} P_{klb} R_{de} D_d R_{ijae} + 2S_{ijkl} P_{ija} R_{de} D_d R_{klbe}. \end{aligned}$$

Separating the terms quadratic in P_{abc} from the rest and noticing that most of the rest is a square, we obtain:

$$\begin{aligned}
& \left(\frac{\partial}{\partial t} - \Delta \right) Z_{ab} \\
&= 2R_{acbd}Z_{cd} \\
&+ 2S_{mntu} (S_{ijrs}P_{ija}\nabla_v R_{rstu} - \nabla_v P_{tua} + R_{tuaw}R_{vw}) \\
&\times (S_{klpq}P_{klb}\nabla_v R_{pqmn} - \nabla_v P_{mnb} + R_{mnbx}R_{vx}) \\
&+ 2R_{acbd}S_{ijkl}P_{ijc}P_{kld} + 2P_{acd}P_{bcd} - 4P_{acd}P_{bdc} - 4S_{ijkl}R_{idae}P_{klb}P_{dje} \\
&- 4S_{ijkl}R_{kdbe}P_{ija}P_{dle} + 2S_{ijmn}S_{klpq}(B_{mnpq} - B_{mnqp} + B_{mpnq} - B_{mqnp})P_{ija}P_{klb} \\
&- 2(S_{dekl}R_{diej} + S_{ijde}R_{dkel})P_{ija}P_{klb} - \frac{1}{2t^2}R_{ab}.
\end{aligned}$$

We simplify the terms that are quadratic in P_{abc} :

$$\begin{aligned}
A \doteq & 2R_{acbd}S_{ijkl}P_{ijc}P_{kld} + 2P_{acd}P_{bcd} - 4P_{acd}P_{bdc} - 4S_{ijkl}R_{idae}P_{klb}P_{dje} \\
& + 2S_{ijmn}S_{klpq}(B_{mnpq} - B_{mnqp} + B_{mpnq} - B_{mqnp})P_{ija}P_{klb} \\
& - 4S_{ijkl}R_{kdbe}P_{ija}P_{dle} - 2(S_{dekl}R_{diej} + S_{ijde}R_{dkel})P_{ija}P_{klb} \\
& = 2R_{acbd}S_{ijkl}P_{ijc}P_{kld} + 2P_{acd}P_{bcd} - 4P_{acd}P_{bdc} \\
& - 4S_{ijkl}R_{idae}P_{klb}P_{dje} - 4S_{ijkl}R_{kdbe}P_{ija}P_{dle} + P_{cda}P_{cdb} \\
& + 2S_{ijmn}S_{klpq}(B_{mpnq} - B_{mqnp})P_{ija}P_{klb} - P_{kla}P_{klb} - P_{ija}P_{ijb} \\
& = -2P_{acd}P_{bdc} + 2R_{acbd}S_{ijkl}P_{ijc}P_{kld} - 4S_{ijkl}R_{idae}P_{klb}P_{dje} \\
& - 4S_{ijkl}R_{idbe}P_{kla}P_{dje} + 2R_{iepf}R_{jeqf}S_{ijkl}P_{kla}S_{pqmn}P_{mnb} \\
& - 2R_{ieqf}R_{jepf}S_{ijkl}P_{kla}S_{pqmn}P_{mnb} \\
& = -2P_{acd}P_{bdc} + 2R_{acbd}S_{ijkl}P_{ijc}P_{kld} - 4S_{ijkl}R_{idae}P_{klb}P_{dje} \\
& - 4S_{ijkl}R_{idbe}P_{kla}P_{dje} + 4R_{iepf}R_{jeqf}S_{ijkl}P_{kla}S_{pqmn}P_{mnb}.
\end{aligned}$$

In the above computation we used the following identities. First, the quadratic B in curvature can be rewritten using

$$\begin{aligned}
R_{mnpq}^2 &= R_{mnef}R_{pqef} = (R_{menf} - R_{nemf})(R_{peqf} - R_{qepf}) \\
&= B_{mnpq} - B_{mnqp} - B_{nmpq} + B_{nmqp} \\
&= 2(B_{mnpq} - B_{mnqp}).
\end{aligned}$$

and

$$\begin{aligned}
& 2S_{ijmn}S_{klpq}(B_{mpnq} - B_{mqnp})P_{ija}P_{klb} \\
&= 2S_{ijmn}S_{klpq}(R_{mepf}R_{neqf} - R_{meqf}R_{nepf})P_{ija}P_{klb} \\
&= 2S_{ijkl}S_{mnpq}R_{iepf}R_{jeqf}P_{kla}P_{mnb} - 2S_{ijkl}S_{mnpq}R_{ieqf}R_{jepf}P_{kla}P_{mnb}.
\end{aligned}$$

Second, we have the identities

$$\begin{aligned}
2S_{dekl}R_{diej}P_{ija} &= P_{kla} \\
2S_{ijde}R_{dkel}P_{klb} &= P_{ijb},
\end{aligned}$$

where the first follows from

$$S_{dekl} (R_{diej} - R_{djei}) P_{ija} = S_{dekl} R_{deij} P_{ija} = P_{kla}$$

which is obtained from

$$S_{dekl} R_{diej} P_{ija} = S_{dekl} (R_{deij} + R_{djei}) P_{ija}$$

and the second is proved similarly. Third, we have the identity

$$\begin{aligned} P_{cda} P_{cdb} &= (P_{adc} + P_{cad}) (P_{bdc} + P_{cbd}) \\ &= 2 (P_{acd} P_{bcd} - P_{acd} P_{bdc}). \end{aligned}$$

Now define $E_{ija} \doteq S_{ijkl} P_{kla}$ to simplify notation further. Then

$$\begin{aligned} A &= -2P_{acd} P_{bdc} + 2R_{acbd} S_{ijkl} P_{ijc} P_{kld} - 4R_{idae} E_{ijb} P_{dje} \\ &\quad - 4R_{idbe} E_{ija} P_{dje} + 4R_{iepf} R_{jeqf} E_{ija} E_{pqb}. \end{aligned}$$

Recall that

$$\begin{aligned} R_{abcd} &= \sum_N Y_{ab}^N Y_{cd}^N \\ P_{abc} &= \sum_N Y_{ab}^N X_c^N \\ M_{ab} &= \sum_N X_a^N X_b^N. \end{aligned}$$

Thus

$$\begin{aligned} A &= -2P_{acd} P_{bdc} + 2R_{acbd} S_{ijkl} P_{ijc} P_{kld} + 4Y_{pf}^N Y_{ah}^N Y_{qf}^M X_h^M E_{pqb} \\ &\quad + 4Y_{id}^N Y_{be}^N Y_{jd}^M X_e^M E_{ija} + 4Y_{id}^N Y_{pf}^N Y_{jd}^M Y_{qf}^M E_{ija} E_{pqb} \\ &= 4 (Y_{id}^N Y_{jd}^M E_{ija} + Y_{ah}^N X_h^M) (Y_{pf}^N Y_{qf}^M E_{pqb} + Y_{be}^N X_e^M) - 4Y_{ah}^N X_h^M Y_{be}^N X_e^M \\ &\quad - 2P_{acd} P_{bdc} + 2R_{acbd} S_{ijkl} P_{ijc} P_{kld} \\ &= 4 (Y_{id}^N Y_{jd}^M E_{ija} + Y_{ah}^N X_h^M) (Y_{pf}^N Y_{qf}^M E_{pqb} + Y_{be}^N X_e^M) - 2R_{acbd} Z_{cd} \\ &\quad - 2R_{acbd} M_{cd} - 2P_{acd} P_{bdc}. \end{aligned}$$

Note that the terms $Y_{id}^N Y_{jd}^M E_{ija}$ and $Y_{pf}^N Y_{qf}^M E_{pqb}$ are both antisymmetric in N and M .

Now the terms on the last line may be rewritten as

$$\begin{aligned} &- 2R_{acbd} M_{cd} - 2P_{acd} P_{bdc} \\ &= -2Y_{ac}^N Y_{bd}^N X_c^M X_d^M - 2Y_{ac}^N X_d^N Y_{bd}^M X_c^M P_{acd} P_{bdc} \\ &= - (Y_{ac}^N X_c^M + Y_{ac}^M X_c^N) (Y_{bd}^N X_d^M + Y_{bd}^M X_d^N) \end{aligned}$$

so that

$$\begin{aligned}
A &= 4 (Y_{id}^N Y_{jd}^M E_{ija} + Y_{ah}^N X_h^M) (Y_{pf}^N Y_{qf}^M E_{pqb} + Y_{be}^N X_e^M) - 2R_{acbd} Z_{cd} \\
&\quad - (Y_{ac}^N X_c^M + Y_{ac}^M X_c^N) (Y_{bd}^N X_d^M + Y_{bd}^M X_d^N) \\
&= (2Y_{id}^N Y_{jd}^M E_{ija} + Y_{ah}^N X_h^M - Y_{ah}^M X_h^N) (2Y_{pf}^N Y_{qf}^M E_{pqb} + Y_{be}^N X_e^M - Y_{be}^M X_e^N) \\
&\quad - 2R_{acbd} Z_{cd},
\end{aligned}$$

where we used

$$Y_{ah}^N X_h^M (Y_{be}^N X_e^M + Y_{be}^M X_e^N) = \frac{1}{2} (Y_{ah}^N X_h^M + Y_{ah}^M X_h^N) (Y_{be}^N X_e^M + Y_{be}^M X_e^N)$$

and

$$Y_{id}^N Y_{jd}^M E_{ija} (Y_{be}^N X_e^M + Y_{be}^M X_e^N) = 0$$

(since $Y_{be}^N X_e^M + Y_{be}^M X_e^N$ is symmetric in N and M).

Therefore the evolution equation for Z_{ab} is:

$$\begin{aligned}
&\left(\frac{\partial}{\partial t} - \Delta \right) Z_{ab} \\
&= 2S_{mntu} (S_{ijrs} P_{ija} \nabla_v R_{rstu} - \nabla_v P_{tua} + R_{tuaw} R_{vw}) \\
&\quad \times (S_{klpq} P_{klb} \nabla_v R_{pqmn} - \nabla_v P_{mnb} + R_{mnbx} R_{vx}) \\
&\quad + (2Y_{id}^N Y_{jd}^M E_{ija} + Y_{ah}^N X_h^M - Y_{ah}^M X_h^N) (2Y_{pf}^N Y_{qf}^M E_{pqb} + Y_{be}^N X_e^M - Y_{be}^M X_e^N) \\
&\quad - \frac{1}{2t^2} R_{ab},
\end{aligned}$$

where $E_{ija} = S_{ijkl} P_{kla}$. We may rewrite this as

$$\begin{aligned}
&\left(\frac{\partial}{\partial t} - \Delta \right) Z_{ab} \\
&= 2S_{mntu} \left(S_{ijrs} P_{ija} \nabla_v R_{rstu} - \nabla_v P_{tua} + R_{tuaw} R_{vw} + \frac{1}{2t} R_{tuav} \right) \\
&\quad \times \left(S_{klpq} P_{klb} \nabla_v R_{pqmn} - \nabla_v P_{mnb} + R_{mnbx} R_{vx} + \frac{1}{2t} R_{mnbv} \right) \\
&\quad + (2Y_{id}^N Y_{jd}^M E_{ija} + Y_{ah}^N X_h^M - Y_{ah}^M X_h^N) (2Y_{pf}^N Y_{qf}^M E_{pqb} + Y_{be}^N X_e^M - Y_{be}^M X_e^N) \\
&\quad - \frac{1}{t} \left(S_{ijrs} P_{ija} \nabla_v R_{rsbv} - \nabla_v P_{bva} + R_{bvaw} R_{vw} + \frac{1}{2t} R_{bvav} \right) \\
&\quad - \frac{1}{t} \left(S_{klpq} P_{klb} \nabla_v R_{pqav} - \nabla_v P_{avb} + R_{avbx} R_{vx} + \frac{1}{2t} R_{avbv} \right)
\end{aligned}$$

Now

$$\begin{aligned}
-\nabla_v P_{bva} + R_{bvaw} R_{vw} + \frac{1}{2t} R_{bvav} &= -\nabla_v (\nabla_b R_{va} - \nabla_v R_{ba}) + R_{bvaw} R_{vw} + \frac{1}{2t} R_{ba} \\
&= \Delta R_{ba} - \nabla_b \nabla_v R_{va} - R_{bw} R_{wa} + R_{bvaw} R_{vw} \\
&\quad + R_{bvaw} R_{vw} + \frac{1}{2t} R_{ba} \\
&= \Delta R_{ba} - \frac{1}{2} \nabla_b \nabla_a R + 2R_{bvaw} R_{vw} - R_{ac} R_{bc} + \frac{1}{2t} R_{ab} \\
&= M_{ab}
\end{aligned}$$

and

$$\nabla_v R_{rsbv} = \nabla_r R_{vsbv} + \nabla_s R_{rvbv} = -\nabla_r R_{sb} + \nabla_s R_{rb} = -P_{rsb}.$$

Hence

$$\begin{aligned}
&\left(\frac{\partial}{\partial t} - \Delta \right) Z_{ab} \\
&= 2S_{mntu} \left(S_{ijrs} P_{ija} \nabla_v R_{rstu} - \nabla_v P_{tua} + R_{tuaw} R_{vw} + \frac{1}{2t} R_{tuav} \right) \\
&\quad \times \left(S_{klpq} P_{klb} \nabla_v R_{pqmn} - \nabla_v P_{mnb} + R_{mnbx} R_{vx} + \frac{1}{2t} R_{mnbv} \right) \\
&\quad + (2Y_{id}^N Y_{jd}^M E_{ija} + Y_{ah}^N X_h^M - Y_{ah}^M X_h^N) (2Y_{pf}^N Y_{qf}^M E_{pqb} + Y_{be}^N X_e^M - Y_{be}^M X_e^N) \\
&\quad - \frac{1}{t} (-S_{ijrs} P_{ija} P_{rsb} + M_{ab}) - \frac{1}{t} (-S_{klpq} P_{klb} P_{pqa} + M_{ab}).
\end{aligned}$$

That is

$$\begin{aligned}
&\left(\frac{\partial}{\partial t} - \Delta \right) Z_{ab} = 2S_{mntu} \left(S_{ijrs} P_{ija} \nabla_v R_{rstu} - \nabla_v P_{tua} + R_{tuaw} R_{vw} + \frac{1}{2t} R_{tuav} \right) \\
&\quad \times \left(S_{klpq} P_{klb} \nabla_v R_{pqmn} - \nabla_v P_{mnb} + R_{mnbx} R_{vx} + \frac{1}{2t} R_{mnbv} \right) \\
&\quad + (2Y_{id}^N Y_{jd}^M E_{ija} + Y_{ah}^N X_h^M - Y_{ah}^M X_h^N) (2Y_{pf}^N Y_{qf}^M E_{pqb} + Y_{be}^N X_e^M - Y_{be}^M X_e^N) \\
&\quad - \frac{2}{t} Z_{ab}.
\end{aligned}$$

This completes the proof of the main theorem.

Now it is interesting to observe the form of the resulting evolution equation when we trace the above computation. Define

$$Z \doteq Z_{aa} = \frac{1}{2} \left(\Delta R + 2|Rc|^2 + \frac{1}{t} R \right) - S_{ijkl} P_{ija} P_{kla}.$$

This trace Harnack quantity is slightly different than (1); however its positivity is of course still a corollary of Hamilton's matrix Harnack estimate. Tracing

equation (2), we have

$$\left(\frac{\partial}{\partial t} - \Delta\right) Z = 2S_{ijkl}K_{mnij}K_{mnkl} + (L_a^{NM})^2 - \frac{2}{t}Z \geq -\frac{2}{t}Z.$$

Note however that an application of the maximum principle does not yield a *direct* proof of the trace Harnack inequality. This is because in order for L_a^{NM} , which appears on the rhs, to be well-defined, we need the matrix Harnack quantity to be nonnegative!

Acknowledgment. We would like to express our gratitude to Richard Hamilton for suggesting to us the method of proof given in this paper.

References

- [C] H.-D. Cao, *On Harnack's inequalities for the Kähler-Ricci flow*, Invent. math. **109** (1992) 247–263.
- [CC] B. Chow and S.C. Chu, *A geometric interpretation of Hamilton's Harnack inequality for the Ricci flow*, Math. Res. Lett. **2** (1995) 701–718.
- [H0] R. S. Hamilton, *The Ricci flow on surfaces*, Contemp. Math. **71** (1988) 237–261.
- [H1] R. S. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Diff. Geom. **17** (1982), 255–306.
- [H2] R. S. Hamilton, *The Harnack estimate for the Ricci flow*, J. Diff. Geom. **37** (1993) 225–243.
- [H3] R. S. Hamilton, *The formation of singularities in the Ricci flow*, in Surveys in Differential Geometry **2** (1995) International Press, 7–136.
- [LY] P. Li and S.-T. Yau, *On the parabolic kernel of the Schrödinger operator*, Acta. Math. **156** (1986) 153–201.